

# The effects of cryotherapy on quadriceps electromyographic activity and isometric strength in patient in the early phases following knee surgery

Journal of Orthopaedic Surgery  
27(1) 1–8

© The Author(s) 2019

Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/2309499019831454  
journals.sagepub.com/home/osj



William A Loro<sup>1</sup>, Mark D Thelen<sup>2</sup>, Michael D Rosenthal<sup>3</sup>,  
Paul D Stoneman<sup>4</sup> and Michael D Ross<sup>5</sup>

## Abstract

**Purpose:** To determine the effects of cryotherapy on quadriceps electromyographic (EMG) activity and isometric strength in early postoperative knee surgery patients. **Methods:** Twenty-two volunteers with recent knee surgeries were included. EMG readings of the vastus medialis (VM), rectus femoris (RF), and vastus lateralis (VL) from the surgical leg were collected during a maximal voluntary quadriceps setting (QS) activity. Maximum isometric knee extension force measurements were also recorded. Subjects were randomly assigned to receive an ice bag or a sham room-temperature bag to the front of their postsurgical knee for 20 min. After treatment, the subjects repeated the above mentioned maximum QS and isometric knee extension force measurements. The subjects returned 24 h later to conduct the same test protocol but received the treatment (ice or sham) not applied during their first test session. **Results:** A 38% increase in VM EMG activity during QS and a 30% increase in maximum isometric knee extension strength were found after cryotherapy treatment. No significant differences were found in RF or VL EMG activity during QS after cryotherapy. No significant differences were found in any measurements after the sham treatment. **Conclusion:** Clinicians should consider applying ice to knee joints prior to exercise for patients following knee surgery with inhibited quadriceps.

## Keywords

electromyography, ice, quadriceps inhibition, rehabilitation

Date received: 1 April 2018; accepted: 13 January 2019

## Introduction

Arthrogenic muscle inhibition of the quadriceps is commonplace after knee surgery.<sup>1–4</sup> Arthrogenic muscle inhibition is a presynaptic, ongoing reflex inhibition of musculature surrounding a joint after distension or damage to the joint structures.<sup>5</sup> While pain and disuse are considered primary reasons for inhibition, arthrogenic muscle inhibition results from activity from many different joint receptors, which act on inhibitory interneurons synapsing on the motor neuron pool of joint musculature.<sup>5</sup> Mechanoreceptor activity may be most responsible for muscle inhibition. Quadriceps inhibition may occur from damage to articular mechanoreceptors which causes abnormal joint

<sup>1</sup> 1-1-1 Physical Therapy, Somers Point, NJ, USA

<sup>2</sup> US Army-Baylor University Doctoral Program in Physical Therapy, Fort Sam Houston, TX, USA

<sup>3</sup> Department of Physical Therapy, San Diego State University, San Diego, CA, USA

<sup>4</sup> Department of Physical Therapy, Rocky Mountain University of Health Professions, Provo, UT, USA

<sup>5</sup> Department of Physical Therapy, Daemen College, Amherst, NY, USA

## Corresponding author:

William A. Loro, 1-1-1 Physical Therapy, 32 W Maryland Ave, Somers Point, NJ 08244, USA.

Email: lorowal4@gmail.com



afferent input that decreases excitability of the spinal neurons controlling quadriceps activity.<sup>6</sup>

Arthrogenic muscle inhibition is a limiting factor in joint rehabilitation, because it delays active exercise necessary for healing.<sup>5</sup> Reactivating patients' inhibited quadriceps remains a primary objective in the early stages of postoperative knee surgery rehabilitation. Methods of reactivating inhibited quadriceps include volitional exercise, the use of electrical stimulation, and biofeedback.<sup>7-11</sup> Cryotherapy may have a role in reactivating inhibited quadriceps. Hopkins et al.<sup>9</sup> observed that cryotherapy and transcutaneous electric neuromuscular stimulation decreased the inhibition of the vastus medialis (VM), and cryotherapy further facilitated the quadriceps motor neuron pool in artificially effused knees. The cryotherapy group experienced greater "disinhibition" or activation of the VM than the transcutaneous electric neuromuscular stimulation group with longer lasting effects.<sup>9</sup> Disinhibition is defined as "a return of some measure of recruitment to baseline or pre-injury levels."<sup>12</sup> Hopkins<sup>12</sup> later studied the effects of cryotherapy again in artificially effused knees. Consistent with earlier work, he observed that joint cryotherapy applied after knee effusion resulted in disinhibition of the quadriceps.<sup>12</sup>

There is no consensus regarding the effects of cryotherapy on voluntary motor activity. Although the majority of authors have reported increases,<sup>7,9,12-17</sup> others have reported decreases<sup>9,18-20</sup> or no change<sup>8,21,22</sup> in muscle activity or force after muscle or joint cooling.<sup>23</sup> Differences in the site (muscle vs joint) of ice application, the type of measurement used, and the time that the measurement was taken (during or after cryotherapy treatment) contribute to the inconsistent results reported in the literature.<sup>23</sup> The use of ice or cryotherapy to assist with pain and inflammation is routine for knee surgery patients.<sup>21,22,24</sup> Yet, the effects of cryotherapy on inhibited quadriceps activity and strength in early postoperative knee patients have not been studied. We hypothesized that cryotherapy applied to the knee joint of early postoperative knee surgery patients would lead to an increase in quadriceps surface electromyographic (EMG) activity and isometric quadriceps strength.

## Materials and methods

### Subjects

Twenty-two (15 males and 7 females) Department of Defense beneficiaries (mean age =  $22.2 \pm 5.4$  years) who had recent knee surgery (mean time of testing following surgery =  $4.3 \pm 2.1$  days) volunteered to participate in this study. Subjects were eligible to participate if they had surgery within 10 days of the completion of data collection. Eligible surgical procedures included isolated anterior cruciate ligament reconstruction (ACL-R); ACL-R with meniscectomy; ACL-R with associated ligamentous and/or soft tissue injury such as the medial collateral ligament,

lateral collateral ligament, and menisci not requiring surgical repair; and arthroscopic knee surgeries to include meniscectomy and/or meniscal debridement, plica excision, and patellar shaving. Subjects were excluded if they had the following: ligamentous repairs other than ACL-R, surgery with meniscal repair or allograft, osteochondral procedures, and nonsurgical knee pathologies.

Of the 22 subjects, 15 had ACL-R surgeries using the semitendinosus and gracilis in single or double-bundled fashion (9) or bone-patellar-tendon-bone (6) as a graft source. The remaining seven subjects had arthroscopic knee surgeries involving meniscal debridement. One subject with arthroscopic meniscal debridement also had a lateral retinacular release.

This study was approved by the Keller Army Community Hospital Institutional Review Board at West Point, New York. All subjects signed informed consent forms before participating, and the rights of all subjects were protected.

### Design

This was a two-by-two repeated-measures design, with each subject tested to assess their response to a type of treatment (ice or no ice/sham) over time (pretreatment and posttreatment). The subjects participated in two separate data collection sessions randomly to evaluate the effects of ice and no ice/sham treatment on their quadriceps surface EMG activity and isometric knee extension strength. Statistical analysis was done using repeated-measure analysis of variance (ANOVA) comparing the pretreatment and posttreatment means of both quadriceps EMG activity and isometric strength with SPSS 11.5 for Windows (SPSS Corp. Chicago, Illinois, USA). Significance level was set at  $p < 0.05$ .

### Quadriceps EMG recording and data analysis

The surface EMG measurements were collected using the Biopac MP100WSW system (BIOPAC Systems, Inc., Goleta, California, USA) and Bagnoli EMG System (Delsys Inc., Boston, Massachusetts, USA). The electrode sites were shaved and scrubbed with rubbing alcohol to prepare the skin.<sup>25,26</sup> Single differential Ag-AgCl bar electrodes (DE-2.1; Delsys Inc.) were positioned using an electrolyte gel and adhesive skin interfaces at a fixed interelectrode distance of 1 cm.<sup>27</sup> The reference electrode was attached to the medial malleolus of the nonsurgical leg. EMG data were collected for the following muscles of the surgical leg: VM (two finger breadths proximal to the superomedial border of the patella along the assumed line of the fibers), rectus femoris (RF; midway along a line between the anterior superior iliac spine and the superior pole of the patella), and vastus lateralis (VL; four finger breadths proximal to the superolateral border of the patella along the assumed line of the fibers).<sup>1</sup> The three sites were

marked with a skin marker prior to electrode placement to duplicate placement during the second data collection session. The electrodes stayed in place for the duration of the test session. Pilot testing revealed that skin surface temperature measurements at the EMG electrode sites did not change during and after ice bag application, which was consistent with comparable studies.<sup>9,12</sup> Therefore, skin surface temperature was not monitored during testing.

The EMG sampling rate was 2000 Hz. The EMG signal was amplified with an overall gain of 1000 and digitized using Acqknowledge 3.7.2 software (BIOPAC Systems, Inc). Bandpass (20–500 Hz) and notch filters (60 Hz) were used. For testing of quadriceps muscle setting, the EMG signal was collected for 5 s during a maximal voluntary isometric contraction (MVIC), whereas the knee was fully extended (isometric action of muscles monitored). The root-mean-square (RMS) for three trials during the middle 3 s (first and last seconds discarded) of the 5-s MVIC was calculated for each muscle. The greatest value of the three consecutive attempts was then calculated for analysis.<sup>27</sup>

### Isometric quadriceps strength measurement and data analysis

Isometric quadriceps strength was measured with a Lafayette Manual Muscle Test System (NEXGEN Ergonomics Inc., Montreal, Quebec, Canada) hand-held dynamometer. Interrater reliability is good to high for measurements of knee extensor force obtained by handheld dynamometry.<sup>28,29</sup> The peak force measurement of an MVIC while attempting knee extension with the knee maintained at 70° flexion was collected for 5 s. Two 5-s trials were averaged for analysis.

### Procedure

The enrolled volunteers reported to outpatient physical therapy to initiate postoperative knee rehabilitation per pre-established rehab protocols. As soon as the subjects acquired 70° of flexion in their surgical knee, not to exceed 10 days postsurgery, they participated in two separate 45-min data collection sessions. The separate sessions took place 24 h apart inside a quiet laboratory. The order of the testing was randomized with 14 of the 22 subjects receiving cryotherapy treatment during the first test session.

Subjects were seated with both knees fully extended on a padded table with their backs supported against a wall. The subjects' knee girth measurements (in centimeter) were taken at the superior margin of the patella at the beginning of each session. Pain levels were recorded via a 10-point visual analog scale<sup>30</sup> to specifically evaluate subjective complaints after knee surgery. A practice period was held for all subjects at the beginning of each data collection session. This time was used to familiarize the subjects with the testing protocol and to practice maximal



**Figure 1.** Surface electromyographic measurements of the vastus medialis, rectus femoris, and vastus lateralis during quadriceps setting.



**Figure 2.** Maximum voluntary isometric contraction with the knee at 70° flexion.

voluntary quadriceps setting (QS) and isometric knee extensions (three to five practice trials).

EMG measurements of the VM, RF, and VL from the surgical leg were collected during three trials of QS (Figure 1). Verbal commands of “ready,” “push,” and “stop” were given to the subjects. In performing the QS exercises, subjects were instructed to push the back of their surgical knee as hard as possible into the padded table that was under their knee until told to stop. Each QS trial lasted 5 s with 30 s rest between each trial. The subjects were instructed to relax while the data were being stored. After the three trials of QS, the subjects were asked to rate their pain on 10-point visual analog scale.<sup>30</sup> The subjects then sat at the edge of the padded table with their surgical leg in 70° of knee flexion (verified by a goniometer). A handheld dynamometer was placed over the anterior aspect of the subjects' shins, just proximal to the ankle joint (Figure 2). Again



**Figure 3.** Setup of treatment application.

using the verbal commands of “ready,” “push,” and “stop,” the subjects were instructed to extend their knee with maximal effort for 5 s with 30 s rest between two trials. The subjects were then asked to rate their pain on a 10-point visual analog scale.<sup>30</sup>

The subjects then received either a 1.5 L crushed ice bag or a room-temperature bag (sham treatment—kitty litter) with the same consistency and weight as an ice bag to the front part of their postsurgical knee for 20 min. The bags were secured to the knee using an elastic ace wrap (Figure 3), which concealed the type of treatment to the blinded tester. Immediately after the cryotherapy or sham treatment, the abovementioned QS and isometric quadriceps strength testing was repeated using the same procedure as pretreatment. The subjects returned to the lab 24 h later to conduct the same test protocol with the treatment (ice or sham) that was not applied during the first test session. Between test sessions, the subjects continued their assigned home exercise programs (per preestablished rehabilitation protocols) consisting primarily of heel slides, QS, straight-leg raises, and calf pumps.

The subjects were blinded to the results of their measurements. The same tester collected all EMG measurements. The tester of the isometric strength testing using the handheld dynamometer was a different individual blinded to the type of treatment. After the second test session, the subjects were discharged from the study and resumed postoperative knee surgery rehabilitation.

### Statistical analysis

A two-by-two repeated-measure analyses of variance (ANOVA) was performed to compare prequadriceps and postquadriceps EMG activity and isometric strength. The within-subject factors were treatment (two levels: ice or no ice/sham treatment) and time (two levels: pretreatment and posttreatment). Paired *t*-tests were computed for knee girth

**Table 1.** Electromyographic activity during maximum quadriceps setting.<sup>a</sup>

	VM	RF	VL
Pre-ice	17.24 (16.63)	12.51 (13.79)	10.81 (12.57)
Post-ice	23.76 (24.95) <sup>b</sup>	13.65 (14.46)	10.29 (11.65)
Pre-sham	19.85 (19.84)	16.06 (20.14)	10.26 (6.58)
Post-sham	21.06 (24.10)	15.04 (24.23)	9.31 (6.53)

VM: vastus medialis, RF: rectus femoris; VL: vastus lateralis.

<sup>a</sup>Values are expressed in mean (standard deviation) are root-mean-square in millivolts.

<sup>b</sup>37.82% increase from pre-ice assessment ( $p \leq 0.024$ ).

**Table 2.** Isometric quadriceps peak force.<sup>a</sup>

	Force (kg)
Pre-ice treatment	11.05 (3.76)
Post-ice treatment	14.35 (4.62) <sup>b</sup>
Pre-sham treatment	11.25 (4.11)
Post-sham treatment	11.84 (4.36)

<sup>a</sup>Values are expressed in mean (standard deviation).

<sup>b</sup>29.86% increase from pre-ice assessment ( $p \leq 0.0001$ ).

and the Wilcoxon signed-ranks test was used for visual analog scale pain scores. Wilcoxon signed-ranks test was used for pain data. A statistical significance level was set at  $p < 0.05$ .

### Results

Means and standard deviations for quadriceps EMG activity and isometric strength are provided in Tables 1 and 2. VM EMG activity increased significantly after cryotherapy treatment ( $F_{1, 21} = 5.922$ ;  $p \leq 0.024$ ). There was no significant difference in VM EMG activity between pre-sham and post-sham treatment ( $F_{1, 21} = 0.047$ ;  $p = 0.530$ ). Likewise, no significant differences were found in RF and vastus lateralis (VL) EMG activity between precryotherapy and postcryotherapy (RF:  $F_{1, 21} = 1.864$ ;  $p = 0.187$ ; VL:  $F_{1, 21} = 0.186$ ;  $p = 0.670$ ) or pre-sham and post-sham treatments (RF:  $F_{1, 21} = 0.284$ ;  $p = 0.600$ ; VL:  $F_{1, 21} = 2.528$ ;  $p = 0.127$ ).

Quadriceps isometric knee extension strength increased significantly after cryotherapy treatment ( $F_{1, 21} = 31.097$ ;  $p \leq 0.0001$ ). No significant difference in quadriceps isometric knee extension strength was found between pre-sham and post-sham treatment ( $F_{1, 21} = 3.743$ ;  $p = 0.067$ ). With respect to knee girth, the difference between involved and noninvolved lower extremities for the two testing days was not significantly different (mean differences between involved and noninvolved lower extremities of  $3.3 \pm 1.5$  cm for testing session 1 and  $3.5 \pm 1.4$  cm for testing session 2). There were also no significant differences in pain levels during each test session or between test days (Table 3).

**Table 3.** Pain levels that were recorded via a 10-point visual analog scale.

	Cryotherapy session	Sham session
Initial pain level	2.50 (1.68)	2.50 (1.54)
Pain level after QS: pretreatment	2.86 (1.58)	2.86 (1.91)
Pain level after isometric quadriceps force assessment: pretreatment	4.23 (2.51)	4.14 (2.36)
Pain level after QS: posttreatment	2.55 (1.44)	2.77 (1.72)
Pain level after isometric quadriceps force assessment: posttreatment	3.95 (2.32)	4.05 (2.65)

QS: quadriceps setting.

## Discussion

The results of this study support our hypothesis stating cryotherapy applied to the knee joint of early postoperative knee surgery patients leads to an increase in quadriceps surface EMG activity and isometric quadriceps strength. Consistent with previous studies,<sup>9,12</sup> we observed that knee joint cryotherapy actually had a disinhibitory effect on the inhibited quadriceps. While Hopkins et al.<sup>9,12</sup> used an experimental knee joint effusion model, our study included patients in the early postoperative phase. The reason why knee joint cryotherapy disinhibits the quadriceps is not fully understood. Arthrogenic muscle inhibition is associated with afferent stimuli from the knee joint<sup>31</sup> and occurs when sensory stimuli impede the voluntary activation of the muscle.<sup>32</sup> Certain sensory stimuli can block other afferent sensory stimuli in the spinal cord. Cutaneous sensory nerve stimulation can increase motor neuron excitability.<sup>9,31</sup> Cryotherapy might reduce quadriceps inhibition by preventing the activation of inhibitory synapses and/or by increasing the excitability of the anterior horn cells.<sup>31</sup> Cryotherapy may result in less supraspinal control over reflexive activity and more recruitment within the quadriceps motor neuron pool.<sup>12</sup>

While pain and effusion may have a role in quadriceps inhibition in postoperative knee patients, we observed no difference in knee girth measurement between test days or for pain level during each test session (pretreatment vs. posttreatment) or between test days. Previous studies have demonstrated, using the artificial knee joint effusion model, that arthrogenic muscle inhibition can occur in the absence of pain.<sup>9,33</sup> Based on our findings, we are in agreement with Hopkins et al.<sup>9,22</sup> that cryotherapy has a direct effect on diminishing arthrogenic muscle inhibition and not just pain.

The use of cryotherapy to specifically diminish arthrogenic muscle inhibition or disinhibit muscles is not widely studied or commonly practiced among rehabilitation professionals. During rehabilitation, cryotherapy is most commonly used after active exercise because of its well-documented local physiological responses of decreased temperature, metabolism, inflammation, circulation, pain,

and muscle spasm.<sup>34</sup> However, using *joint* cryotherapy to assist in the reactivation of inhibited muscles is an unconventional, yet promising, concept in clinical application.<sup>9,12,23</sup> The opportunity to assess patients in the early phase following knee surgery in this study is clinically relevant. Our findings, combined with the work of Hopkins et al.,<sup>9,12,23</sup> are pertinent to actual patients seen in rehabilitation settings.

Sixty-eight percent (15 of 22) of the subjects received ACL-R surgeries. The scarcity of existing literature regarding the early postoperative strength testing of patients following ACL-R suggests that maximal isometric quadriceps testing in these patients may be unsafe and might compromise the surgical grafts. But according to Beynonn et al.,<sup>35–37</sup> isometric quadriceps strength measurement is safely attainable for patients following ACL-R, if the patients are tested between 60° and 90° of knee flexion. Based on these findings, we felt testing at 70° knee flexion was safe for our subjects. All 22 enrolled subjects completed data collection without adverse events or effects.

The VM was the only portion of the quadriceps that we found a difference in EMG activity after cryotherapy. This difference may be due to the VM being the most affected portion of quadriceps from arthrogenic muscle inhibition. While common belief and clinical observation suggest that the VM is more inhibited than the other heads of the quadriceps after joint injury,<sup>5,31</sup> recent data indicate uniform weakness, atrophy, and deactivation of the quadriceps after joint injury.<sup>38,39</sup>

QS is a primary exercise assigned to patients with inhibited quadriceps. Generally in most rehabilitation settings, QS is done by forcefully contracting the quadriceps with the knee in maximum extension. It has been suggested that QS should be performed with the knee in a position of some flexion (approximately 30°) to avoid inhibition of the muscle.<sup>32,40</sup> The decision to conduct QS during this experiment in full extension was based on the authors' intent to keep the testing protocol consistent with general clinical practice. Each subject's postsurgical knee range of motion was measured and verified to be at least 0–70 prior to each test session. However, each subject's postsurgical knee extension was not compared to their nonsurgical knee range of motion. Therefore, symmetrical bilateral knee extension, and potentially hyperextension, was not verified at the time of testing. Lacking knee hyperextension (if hyperextension is normal) may contribute to decreased quadriceps activation during quadriceps setting.<sup>41,42</sup> Whether or not the subjects' surgical knee range of motion would have changed between pretreatment and posttreatment and between testing days is questionable. The time it takes for postsurgical knee patients to reacquire knee hyperextension, if normal, is patient specific and highly variable. After a subsequent review of the subjects' medical records, we found no documentation of genu recurvatum or knee hyperextension in any of the subjects' bilateral knees. Retrospectively, the

subjects' nonsurgical knee range of motion should have been verified at the time of testing.

With a repeated-measures design with testing sessions separated by 24 h, it is fairly safe to assume that subject characteristics (i.e. pain, motivation) and testing conditions (i.e. time of day, room temperature) remained constant throughout the course of this experiment. Therefore, differences seen among treatment conditions are more likely to reflect treatment effects and not variability between subjects.<sup>43</sup> Of the quadriceps inhibition and EMG studies reviewed, gender differences among the subjects either did not exist or were not mentioned.<sup>1,9,12,26,40,44–50</sup> One disadvantage of a repeated-measure design was the potential for practice effects.<sup>43</sup> The additional data collection day to evaluate the effects of no ice/sham treatment helped to minimize practice effects in this study. The order of testing, QS followed by isometric quadriceps test at 70°, was not randomized. We assumed that isometric quadriceps extension testing would be uncomfortable for early postoperative knee patients, which might have resulted in subjects' effort being affected during ensuing QS.

Relatively high standard deviations were observed in the EMG activity scores during QS. This finding is consistent with similar EMG studies.<sup>12</sup> Variance in subjects' EMG scores in our study was not surprising, because the subjects were not expected to demonstrate equal levels of quadriceps inhibition after knee surgery. We observed significant differences only in VM EMG activity and overall isometric quadriceps strength after cryotherapy. Our relatively sample size of only 22 subjects may have contributed to not finding a difference in the RF and VL after cryotherapy and in all muscles after the sham treatment. Our clinically meaningful effect size was a 15% increase in both quadriceps EMG activity and isometric strength, which was exceeded in this study for VM EMG activity and overall isometric quadriceps strength after cryotherapy. A few of the measurements approached significance including RF EMG activity postcryotherapy ( $p = 0.187$ ), VL EMG activity post-sham ( $p = 0.127$ ), and isometric strength post-sham ( $p = 0.067$ ). Yet, all these measurements showed small differences that are not considered clinically meaningful.

Our results cannot be generalized to all populations, because 19 of the 22 subjects who participated in this study were healthy young military cadets. The remaining subjects (two males, ages 29 and 41; one female, age 33) were active, healthy military members. The changes seen in VM EMG activity and isometric knee extension strength after icing the postoperative knee joints of individuals with quadriceps inhibition should be confirmed in other patient populations. Also, we did not investigate the parameters of how long to ice the joint nor how long the disinhibitory effects of cryotherapy last. Our sample was also somewhat heterogeneous. While all of the subjects in this study had pain and joint effusion, the degree of pain and joint effusion may vary among patients and the different surgical

procedures which would influence muscle recruitment and strength. Future research should include randomized clinical trials comparing outcomes of patients following knee surgery who receive cryotherapy to their knee joints prior to active exercise to matched control groups. Also, the selective inhibition of the VM in patients following knee surgery and the effects of cryotherapy solely on VM inhibition and reactivation warrant further investigation.

## Conclusion

This study examined the immediate effects of cryotherapy on quadriceps EMG activity and isometric strength in patients following knee surgery in the early postoperative phase. Our results indicate that cryotherapy applied to the knee joint leads to an increase in VM EMG activity and quadriceps isometric strength. Rehabilitation clinicians should consider the use of ice prior to active exercise for knee surgery patients with inhibited quadriceps.

## Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## References

1. McHugh MP, Tyler TF, Stephen JN, et al. Electromyographic analysis of quadriceps fatigue after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 2001; 31(1): 25–32.
2. Shakespeare DT, Stokes M, Sherman KP, et al. Reflex inhibition of the quadriceps after meniscectomy: lack of association with pain. *Clin Physiol* 1985; 5: 137–144.
3. Snyder-Mackler L, De Luca PF, Williams PR, et al. Reflex inhibition of the quadriceps femoris muscle after injury or reconstruction of the anterior cruciate ligament. *J Bone Joint Surg Am* 1994; 76(4): 555–560.
4. Snyder-Mackler L, Delitto A, Bailey SL, et al. Strength of the quadriceps femoris muscle and functional recovery after reconstruction of the anterior cruciate ligament. A prospective, randomized clinical trial of electrical stimulation. *J Bone Joint Surg Am* 1995; 77(8): 1166–1173.
5. Hopkins JT and Ingersoll CD. Arthrogenic muscle inhibition: a limiting factor in joint rehabilitation. *J Sport Rehabil* 2000; 9: 135–159.
6. Hurley MV, Jones DW, and Newham DJ. Arthrogenic quadriceps inhibition and rehabilitation of patients with extensive traumatic knee injuries. *Clin Sci* 1994; 86: 305–310.
7. Delitto A, Rose SJ, McKowen JM, et al. Electrical stimulation versus voluntary exercise in strengthening thigh

- musculature after anterior cruciate ligament surgery. *Phys Ther* 1988; 68(5): 660–663.
8. Draper V. Electromyographic biofeedback and recovery of quadriceps femoris muscle function following anterior cruciate ligament reconstruction. *Phys Ther* 1990; 70(1): 11–17.
  9. Hopkins JT, Ingersoll CD, Edwards J, et al. Cryotherapy and transcutaneous electric neuromuscular stimulation decrease arthrogenic muscle inhibition of the vastus medialis after knee joint effusion. *J Athl Train* 2001; 37(1): 25–31.
  10. Lieber RL, Silva PD, and Daniel DM. Equal effectiveness of electrical and volitional strength training for quadriceps femoris muscles after anterior cruciate ligament surgery. *J Ortho Res* 1996; 14(1): 131–138.
  11. Shaw T, Williams MT, and Chipchase LS. Do early quadriceps exercises affect the outcome of ACL reconstruction? A randomized controlled trial. *J Physiother* 2005; 51: 9–17.
  12. Hopkins JT. Knee joint effusion and cryotherapy alter lower chain kinetics and muscle activity. *J Athl Train* 2006; 41(2): 177–184.
  13. Kinugasa R, Yoshida K, and Horii A. The effects of ice application over the vastus medialis on the activity of quadriceps femoris assessed by muscle function magnetic resonance imaging. *J Sports Med Phys Fitness* 2005; 45: 360–364.
  14. Kinugasa R, Yoshida K, Takayuki W, et al. Skin cooling alters the activation patterns of different heads of the quadriceps. *Can J Appl Physiol* 2005; 30(2): 127–139.
  15. McGown HL. Effects of cold application on maximal isometric contraction. *Phys Ther* 1967; 47(3): 185–192.
  16. Oliver RA, Johnson DJ, Wheelhouse WW, et al. Isometric muscle contraction response during recovery from reduced intramuscular temperature. *Arch Phys Med Rehabil* 1979; 60: 126–129.
  17. Sanya AO and Bello AO. Effects of cold application on isometric strength and endurance of quadriceps femoris muscle. *Afr J Med Sci* 1999; 28(3): 195–198.
  18. Hatzel BM and Kaminski TW. The effects of ice immersion on concentric and eccentric isokinetic muscle performance in the ankle. *Isokinet Exerc Sci* 2000; 8: 103–107.
  19. Maattacola CG and Perrin DH. Effects of cold water submersion on isokinetic strength of the plantar flexors. *Isokinet Exerc Sci* 1993; 3: 152–154.
  20. Ruiz DH, Myrer JW, Durrant E, et al. Cryotherapy and sequential exercise bouts following cryotherapy on concentric and eccentric strength in the quadriceps. *J Athl Train* 1993; 28(4): 320–323.
  21. Konrath GA, Lock T, Goitz HT, et al. The use of cold therapy after anterior cruciate ligament reconstruction. *Am J Sports Med* 1996; 24(5): 629–633.
  22. Lessard LA, Scudds RA, Amendola A, et al. The efficacy of cryotherapy following arthroscopic knee surgery. *J Ortho Sports Phys Ther* 1997; 26(1): 14–22.
  23. Hopkins JT and Stencil RS. Ankle cryotherapy facilitates soleus function. *J Ortho Sports Phys Ther* 2002; 32(12): 622–627.
  24. Raynor MC, Pietrobon R, Guller U, et al. Cryotherapy after ACL reconstruction. *J Knee Surg* 2005; 18(2): 123–129.
  25. Konrad P. A practical introduction to kinesiological electromyography. In: Noraxon INC. USA (ed) *The ABC of EMG*. Scottsdale, AZ: Noraxon INC, 2005, pp. 10–21.
  26. Soderberg GL and Cook TM. An electromyographic analysis of quadriceps femoris muscle setting and straight leg raising. *Phys Ther* 1983; 63(9): 1434–1438.
  27. De Luca CJ. The use of surface electromyography in biomechanics. *J Appl Biomech* 1997; 13(2): 135–163.
  28. Bohannon RW. Test-retest reliability of hand-held dynamometry during a single session of strength assessment. *Phys Ther* 1986; 66(2): 206–209.
  29. Dunn JC and Iversen MD. Interrater reliability of knee muscle forces obtained by hand-held dynamometer from elderly subjects with degenerative back pain. *J Geriatr Phys Ther* 2003; 26(3): 23–29.
  30. Flandry F, Hunt JP, Terry G, et al. Analysis of subjective knee complaints using visual analog scales. *Am J Sports Med* 1991; 19(2): 112–118.
  31. Stokes M and Young A. The contribution of reflex inhibition to arthrogenous muscle weakness. *Clin Sci* 1984; 67: 7–14.
  32. Morrissey MC. Reflex inhibition of thigh muscles in knee injury: causes and treatment. *Sports Med* 1989; 7: 263–276.
  33. Spencer JD, Hayes KC, and Alexander IJ. Knee joint effusion and quadriceps reflex inhibition in man. *Arch Phys Med Rehabil* 1984; 65: 171–177.
  34. Knight KL. *Cryotherapy in sport injury management*. Champaign, IL: Human Kinetics, 1995.
  35. Beynnon BD and Fleming BC. Anterior cruciate ligament strain in-vivo: a review of previous work. *J Biomech* 1998; 31: 519–525.
  36. Beynnon BD, Fleming BC, Johnson RJ, et al. Anterior cruciate ligament strain behavior during rehabilitation exercises in vivo. *Am J Sports Med* 1995; 23(1): 24–35.
  37. Beynnon BD, Johnson RJ, Fleming BC, et al. The strain behavior of the anterior cruciate ligament during squatting and active flexion-extension: a comparison of an open and a closed kinetic chain exercise. *Am J Sports Med* 1997; 25(6): 823–829.
  38. Akima H and Furukawa T. Atrophy of thigh muscles after meniscal lesions and arthroscopic partial meniscectomy. *Knee Surg Sports Traumatol Arthrosc* 2005; 13(8): 632–637.
  39. Williams GN, Buchanan TS, Barrance PJ, et al. Quadriceps weakness, atrophy, and activation failure in predicted noncopers after anterior cruciate ligament injury. *Am J Sports Med* 2005; 33(3): 402–407.
  40. Schulthies SS, Ricard MD, Alexander KJ, et al. An electromyographic investigation of 4 elastic-tubing closed kinetic chain exercises after anterior cruciate ligament reconstruction. *J Athl Train* 1998; 33(4): 328–335.
  41. Grood ES, Suntay WJ, Noyes FR, et al. Biomechanics of the knee-extension exercise. Effect of cutting the anterior cruciate ligament. *J Bone Joint Surg Am* 1984; 66(5): 725–734.
  42. Shelbourne KD and Klotz C. What I have learned about the ACL: utilizing a progressive rehabilitation scheme to achieve total knee symmetry after anterior cruciate ligament reconstruction. *J Ortho Sci* 2006; 11(3): 318–325.

43. Portney LG and Watkins MP. *Foundations of clinical research: applications to practice*, 2nd ed. Upper Saddle River, NJ: Prentice-Hall, 2000.
44. Beutler AI, Cooper LW, Kirkendall DT, et al. Electromyographic analysis of single-leg, closed chain exercises: implications for rehabilitation after anterior cruciate ligament construction. *J Athl Train* 2002; 37(1): 13–18.
45. Gyzlo SM, Patek RM, Pink M, et al. Electromyographic analysis of knee rehabilitation exercises. *J Ortho Sports Phys Ther* 1994; 20(1): 36–43.
46. Hanten WP and Schulthies SS. Exercise effect on electromyographic activity of the vastus medialis oblique and vastus lateralis muscles. *Phys Ther* 1990; 70(9): 561–565.
47. McHugh MP, Tyler TF, Browne MG, et al. Electromyographic predictors of residual quadriceps muscle weakness after anterior cruciate ligament reconstruction. *Am J Sports Med* 2002; 30(3): 334–339.
48. Soderberg GL, Minor SD, Arnold K, et al. Electromyographic analysis of knee exercises in healthy subjects and in patients with knee pathologies. *Phys Ther* 1987; 67(11): 1691–1696.
49. Stratford P. Electromyography of the quadriceps femoris muscles in subjects with normal knees and acutely effused knees. *Phys Ther* 1981; 62(3): 279–283.
50. Tho KS, Nemeth G, Lamontagne M, et al. Electromyographic analysis of muscle fatigue in anterior cruciate ligament deficient knees. *Clin Orthop and Relat Res* 1997; 340: 142–151.